




# Your Treatment Discussion Guide

When you think about your current treatment plan for chronic immune thrombocytopenia, have you been wondering if PROMACTA® (eltrombopag) is right for you?

There are a number of factors that go into determining the best course of treatment that your doctor may want to discuss with you, including daily habits, symptoms, and other aspects of your medical condition. PROMACTA may be an option for you and your doctor to consider if you have questions about your current treatment. Together you can determine a treatment plan that's best fit for you.

PROMACTA is a prescription medicine used to treat adults and children 1 year or older with low blood platelet counts due to chronic immune thrombocytopenia (ITP) when other medicines to treat your ITP or surgery to remove the spleen have not worked well enough. PROMACTA is used to try and raise platelet counts in order to lower your risk for bleeding. PROMACTA is not used to make platelet counts normal.

The conversation starters below may be helpful in guiding a discussion about PROMACTA with your doctor.

IF YOU'D LIKE TO DISCUSS...	ASK	SHARE
<p><b>Symptoms or platelet control</b></p> 	<ul style="list-style-type: none"> <li><input type="checkbox"/> Which treatments can help me stabilize my platelet counts?</li> <li><input type="checkbox"/> Why may PROMACTA be right for me?</li> <li><input type="checkbox"/> How long does it take PROMACTA to start working?</li> <li><input type="checkbox"/> Are there limits on how long I can take PROMACTA?</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> How you've felt since your last visit</li> <li><input type="checkbox"/> Any symptoms you're having trouble managing</li> <li><input type="checkbox"/> What you're doing to cope with your symptoms</li> <li><input type="checkbox"/> Any concerns you have about being able to control your symptoms or platelets</li> </ul>
<p><b>The convenience of taking medication</b></p> 	<ul style="list-style-type: none"> <li><input type="checkbox"/> How is PROMACTA given?</li> <li><input type="checkbox"/> How often would I take PROMACTA?</li> <li><input type="checkbox"/> What foods should I avoid when taking PROMACTA?</li> <li><input type="checkbox"/> How would I set up a dosing schedule for PROMACTA that works for me?</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> Any problems you have taking your current medication</li> <li><input type="checkbox"/> How you prefer to take your medication</li> <li><input type="checkbox"/> What your current daily routine is</li> <li><input type="checkbox"/> Any food restrictions you have with your current medication</li> </ul>
<p><b>Side effects</b></p> 	<ul style="list-style-type: none"> <li><input type="checkbox"/> What are the side effects with PROMACTA?</li> <li><input type="checkbox"/> How should I track side effects of PROMACTA?</li> <li><input type="checkbox"/> If I experience side effects, what do I do?</li> <li><input type="checkbox"/> Who could I talk to if I experience side effects with PROMACTA?</li> </ul>	<ul style="list-style-type: none"> <li><input type="checkbox"/> What side effects you're currently experiencing</li> <li><input type="checkbox"/> How you're managing your side effects</li> <li><input type="checkbox"/> Any side effects that are getting worse</li> <li><input type="checkbox"/> How side effects are affecting your daily life</li> </ul>



**TALK TO YOUR DOCTOR TODAY TO SEE IF PROMACTA IS RIGHT FOR YOU.**

Please see Important Safety Information for PROMACTA on pages 2-3. [Click here](http://PROMACTA.com) to visit PROMACTA.com for full Prescribing Information for PROMACTA, including Boxed WARNING, and Medication Guide.



# APPROVED USES AND IMPORTANT SAFETY INFORMATION

## Approved Uses for PROMACTA® (eltrombopag) Tablets

PROMACTA is a prescription medicine used to treat adults and children 1 year and older with low blood platelet counts due to chronic immune thrombocytopenia (ITP) when other medicines to treat your ITP or surgery to remove the spleen have not worked well enough. PROMACTA is used to try to raise platelet counts in order to lower your risk for bleeding.

PROMACTA is not used to make platelet counts normal.

PROMACTA is for treatment of certain people with low platelet counts caused by chronic ITP, chronic hepatitis C virus (HCV), or severe aplastic anemia (SAA), not for a precancerous condition called myelodysplastic syndromes (MDS) or low platelet counts caused by other conditions or diseases.

It is not known if PROMACTA is safe and effective in children with chronic HCV or previously treated SAA, in children younger than 1 year with ITP, or children younger than 2 years when used in combination with standard immunosuppressive therapy as the first treatment for SAA.

## Important Safety Information for PROMACTA® (eltrombopag) Tablets

### What is the most important information I should know about PROMACTA?

PROMACTA can cause serious side effects, including:

#### Liver problems.

PROMACTA may increase your risk of liver problems that may be severe and possibly life-threatening. Your health care provider will do blood tests to check your liver function before you start taking PROMACTA and during treatment. Your health care provider may stop your treatment with PROMACTA if you have changes in your liver function blood tests.

Tell your health care provider right away if you have any of these signs and symptoms of liver problems:

- yellowing of the skin or the whites of the eyes (jaundice)
- unusual darkening of the urine
- unusual tiredness
- right upper stomach area (abdomen) pain
- confusion
- swelling of the stomach area (abdomen)

### What are the possible side effects of PROMACTA?

PROMACTA may cause serious side effects, including:

- **Worsening of a precancerous blood condition to a blood cancer called acute myelogenous leukemia (AML).** PROMACTA is not for treatment of people with a precancerous condition called myelodysplastic syndromes (MDS). If you have MDS and receive PROMACTA, your MDS condition may worsen and become AML. If MDS worsens to become AML, you may die sooner from AML.
- **High platelet counts and higher risk for blood clots.** Your risk of getting a blood clot is increased if your platelet count is too high during treatment with PROMACTA. Your risk of getting a blood clot may also be increased during treatment with PROMACTA if you have normal or low platelet counts. You may have severe problems or die from some forms of blood clots, such as clots that travel to the lungs or that cause heart attacks or strokes. Your health care provider will check your blood platelet counts and change your dose or stop PROMACTA if your platelet counts get too high. Tell your health care provider right away if you have signs and symptoms of a blood clot in the leg such as swelling, pain, or tenderness.

People with chronic liver disease may be at risk for a type of blood clot in the stomach area. Tell your health care provider right away if you have stomach area pain that may be a symptom of this type of blood clot

- **New or worsened cataracts (a clouding of the lens in the eye).** New or worsened cataracts have happened in people taking PROMACTA. Your health care provider will check your eyes before and during your treatment with PROMACTA. Tell your health care provider about any changes in your eyesight while taking PROMACTA

### What should I tell my health care provider before taking PROMACTA?

**Before you take PROMACTA, tell your health care provider about all of your medical conditions, including if you:**

- have liver problems
- have a precancerous condition called MDS or a blood cancer
- have or have had a blood clot
- have a history of cataracts
- have had surgery to remove your spleen (splenectomy)
- have bleeding problems
- are of Asian ancestry (such as Chinese, Japanese, Taiwanese, or Korean). You may need a lower dose of PROMACTA

[Click here to visit PROMACTA.com](https://www.promacta.com) for full Prescribing Information for PROMACTA, including Boxed WARNING, and Medication Guide.



## Important Safety Information for PROMACTA® (eltrombopag) (cont)

- are pregnant or plan to become pregnant. It is not known if PROMACTA will harm an unborn baby. Tell your health care provider if you become pregnant or think you may be pregnant during treatment with PROMACTA. If you are a woman who is able to become pregnant, you must use reliable birth control (contraception) while taking PROMACTA and for at least 7 days after you stop taking PROMACTA. Talk to your health care provider about options of effective birth control methods that may be right for you during this time
- are breastfeeding or plan to breastfeed. You should not breastfeed during treatment with PROMACTA. Talk to your health care provider about the best way to feed your baby during this time

**Tell your health care provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. PROMACTA may affect the way certain medicines work. Certain other medicines may affect the way PROMACTA works.

Especially tell your health care provider if you take:

- certain medicines used to treat high cholesterol, called “statins”
- a blood thinner medicine

Certain medicines may keep PROMACTA from working correctly. Take PROMACTA at least 2 hours before or 4 hours after taking these products:

- antacids used to treat stomach ulcers or heartburn
- multivitamins, mineral supplements, or products that contain iron, calcium, aluminum, magnesium, selenium, and zinc

Ask your health care provider if you are not sure if your medicine is one that is listed above.

Know the medicines you take. Keep a list of them and show it to your health care provider and pharmacist when you get a new medicine.

### What should I avoid while taking PROMACTA?

Avoid situations and medicines that may increase your risk of bleeding.

**The most common side effects of PROMACTA in adults when used to treat chronic immune thrombocytopenia (ITP) are:**

- nausea
- diarrhea
- upper respiratory tract infection (symptoms may include runny nose, stuffy nose, and sneezing)
- vomiting
- urinary tract infection
- pain or swelling (inflammation) in your throat or mouth (oropharyngeal pain and pharyngitis)
- abnormal liver function tests
- muscle aches

**The most common side effects of PROMACTA in children 1 year and older when used to treat chronic ITP are:**

- upper respiratory tract infection (symptoms may include runny nose, stuffy nose, and sneezing)
- pain or swelling (inflammation) in your nose or throat (nasopharyngitis)

Laboratory tests may show abnormal changes to the cells in your bone marrow.

Tell your health care provider about any bruising or bleeding that happens while you take or after you stop taking PROMACTA.

Tell your health care provider if you have any side effect that bothers you or does not go away.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch), or call 1-800-FDA-1088.

**Please see full Prescribing Information, including Boxed WARNING, and Medication Guide.**

**[Click here](http://PROMACTA.com) to visit PROMACTA.com for full Prescribing Information for PROMACTA, including Boxed WARNING, and Medication Guide.**

